

this systematization, the relative probabilities obtained experimentally were normalized to the Z-law, and for concreteness the quantities in the numerators of the fractions were those pertaining to atoms having the larger electron affinity:

$$\varphi = \frac{p(Z_1)}{p(Z_2)} \left| \frac{k_1 Z_1}{k_2 Z_2} \right.$$

In the last column of Table III, the plus sign denotes those experiments in which  $\varphi > 1$ , with the deviation of  $\varphi$  from unity exceeding one rms error; the zero sign denotes experiments in which  $\varphi = 1$  within the limits of errors, and the minus sign denotes experiments with the opposite tendency. Data on the electron affinities of the atoms were taken from the handbook (see [9]). As can be seen from the table, only in five out of 31 cases is the tendency to preferred formation of the mesic atom of the element with the larger electron affinity violated.

One exception is  $\text{Sb}_2\text{O}_3$ , where there are no reliable published data on the electron affinity of antimony, and it is therefore not excluded that the deviation is accidental. Violation takes place for four carbon compounds, which apparently is connected with the very complicated spatial configuration of these molecules.

The measurement procedure and also a detailed discussion of the results will be published later.

<sup>1</sup> E. Fermi and E. Teller, Phys. Rev. 72, 399 (1947).

<sup>2</sup> A. Fafarman and M. H. Shamos, Phys. Rev. 100, 847 (1955).

<sup>3</sup> M. B. Sterns and M. Sterns, Phys. Rev. 100, 847 (1955).

<sup>4</sup> Sens, Swanson, Telegdi, and Yovanovich, Nuovo cimento 7, 536 (1958).

<sup>5</sup> Lathrop, Lundy, Swanson, Telegdi, and Yovanovich, Nuovo cimento 15, 831 (1960).

<sup>6</sup> Astbury, Hattersley, Hussain, Kemp, and Muirhead. Nuovo cimento 18, 1267 (1960).

<sup>7</sup> Echausse, Fillipas, Sutton, Welsh, and Romanowski, Nuovo cimento 24, 666 (1962).

<sup>8</sup> Baijal, Dias, Kaplan, and Pyle, Nuovo cimento 30, 3 (1963).

<sup>9</sup> Spravochnik Énergiya razryva khimicheskikh svyazei, razdel Potentsialy ionizatsii i srodstvo k élektronu (Handbook, Breaking Energies of Chemical Bonds, chapter on Ionization Potentials and Electron Affinity), Acad. Sci. Press, 1962.

## CAPTURE OF NEGATIVE MUONS BY PURE CHROMIUM AND NICKEL ISOTOPES

V. D. BOBROV, V. G. VARLAMOV, Yu. M. GRASHIN, B. A. DOLGOSHEIN, V. G. KIRILLOV-UGRYUMOV, V. S. ROGANOV, A. V. SAMOÏLOV, and S. V. SOMOV

Moscow Engineering-physics Institute

Submitted to JETP editor December 26, 1964

J. Exptl. Theoret. Phys. (U.S.S.R.) 48, 1199 (April, 1965)

THE interaction between a negative muon and a complex nucleus can be approached from two points of view: 1) if the structure of the nucleus is known, the interaction between the muon and the nucleus can help determine the interaction constants; 2) if we know the constants of the interaction of the muon with the nucleus, the muon can be used as an instrument for the study of the structure of the nucleus.

It seems to us that it would be of interest for nuclear physics to obtain data on the capture of muons by nuclei with closed neutron or proton subshells. We used such nuclei in the form of isotopes of Cr with mass numbers 50, 52, 53, and 54, of which  $\text{Cr}^{52}$  has a closed neutron subshell; a closed proton subshell occurs in nickel isotopes with mass numbers 58, 60, and 62. We present below data on the enrichment of these isotopes (in %):

$\text{Cr}^{50}$	$\text{Cr}^{52}$	$\text{Cr}^{53}$	$\text{Cr}^{54}$	$\text{Ni}^{58}$	$\text{Ni}^{60}$	$\text{Ni}^{62}$
90.3	99.5	84.0	78.5	97.9	96.8	89.0

A muon beam from the JINR (Joint Institute of Nuclear Research) synchrocyclotron was used for the experiments.

The total probability of muon capture was determined by measuring the lifetimes of the muon on the K orbit of the atom of the corresponding element. We present the experimental results obtained by processing the corresponding time distributions with a computer:

Nucleus	$\text{Cr}^{50}$	$\text{Cr}^{52}$	$\text{Cr}^{53}$	$\text{Cr}^{54}$	$\text{Ni}^{58}$	$\text{Ni}^{60}$	$\text{Ni}^{62}$
capture rate, $10^5 \text{ sec}^{-1}$	$38.25 \pm 0.50$	$34.52 \pm 0.47$	$32.97 \pm 0.45$	$30.57 \pm 0.42$	$61.10 \pm 1.05$	$55.62 \pm 0.97$	$47.16 \pm 0.95$

A detailed description of the results and of the measurement procedure will be published later.